


### Joseph G. Ouslander, MD

Professor of Geriatric Medicine Senior Advisor to the Dean for Geriatrics Charles E. Schmidt College of Medicine Professor (Courtesy), Christine E. Lynn College of Nursing Florida Atlantic University

Executive Editor, Journal of the American Geriatrics Society





## Disclosures

- Dr. Ouslander is a full-time employee of Florida Atlantic University (FAU) and has received support through FAU for research on INTERACT from the National Institutes of Health, the Centers for Medicare & Medicaid Services, The Commonwealth Fund, the Retirement Research Foundation, PointClickCare, Medline Industries, and Think Research.
- Dr. Ouslander and his wife receive royalties from FAU and Pathway Health for training on and licensing of the INTERACT program.
- Work on funded INTERACT projects is subject to the terms of Conflict of Interest Management plans developed and approved by the FAU Financial Conflict of Interest Committee.

### **FAU** MEDICINE

## COVID-19 in Long-Term Care Facilities: An Update

## **Key Points: The Setting and Population**

- LTCF is a broad term that can include many types of facilities. This presentation focuses on LTCFs that are generally referred to as "skilled nursing facilities", "nursing facilities", and "nursing homes"
- People who reside in these facilities are there for different reasons and differ clinically
  - "Patients" who are there for post-acute care after discharge from the hospital
  - o "Residents" who require long-term care
- LTCF patients and residents are generally at high risk for complications of and mortality from COVID-19
  - o Multiple chronic conditions
  - Advance age






#### Key Points: Presentation of Covid-19 and the Importance of Infection Control

- LTCF patients and residents frequently <u>do not</u> have typical symptoms and signs of COVID-19
  - No symptoms up to 50% or higher
  - Atypical symptoms e.g. low grade temperature elevation; altered mental or functional status; GI symptoms
- LTCF staff may have no symptoms, no fever, and pass multiple screening tests, and still be infected
  - They also may be working multiple jobs at different facilities and be at high risk
  - o They can therefore infect other staff and residents without knowing it
- The only way to prevent infection and further spread of infection is <u>behavior</u> – intensive infection control procedures








## Key Points: Availability of PPE and Testing


- Shortages of PPE persist and will recur in many areas
  - CDC guidelines should be followed to preserve PPE
- Availability of testing is variable and is still hard to get in a timely way except during suspected outbreaks
  - This further highlights the necessity of intensive infection control procedures





## **FAU** MEDICINE





#### **Key Points: Advance Care Planning**

- The mortality rate from COVID-19 will be high in the LTCF population, and clinical deterioration can occur rapidly
- Clinicians and LTCF staff should therefore conduct advance care planning discussions and update advance directives in light of the Coronavirus pandemic
  - The pandemic provides an opportunity to hold discussions that may have been difficult previously and to create COVID specific directives
- Advance Care Planning requires a team approach
  - Ultimately, this requires a trusting relationship between the patient/resident and the team
  - Engage local palliative care and hospice clinicians and teams where available



#### **Key Points: Advance Care Planning**

- Many educational and documentation tools are available
  - Using evidence on prognosis (e.g. <u>www.ePrognosis.com</u>) and simple languaç descriptions of risks and benefits, such as those available in the INTERACT program are helpful
  - $\circ~$  Being clear about the limited meaning of "DNR" is also helpful
  - COVID-19 specific tools are available
    - https://respectingchoices.org/covid-19-resources/
    - https://www.vitaltalk.org/guides/covid-19-communication-skills/
    - <u>https://www.capc.org/toolkits/covid-19-response-resources/</u>
- Documenting and communicating discussions and decisions is critical so that hospital transfers and other interventions are either implemented or withheld based on the patient/resident and family preferences
- Be prepared for patients/residents dying in the facility
  - Check emergency kits and stock with medications for comfort
    - Liquid morphine injectable and oral/sublingual for respiratory distress
    - Lorazepam injectable and oral/sublingual for anxiety/agitation
    - Atropine liquid for secretions



Education on CPR for Residents and Families

### **FAU** MEDICINE

### **COVID-19 in Long-Term Care Facilities: An Update Key Points: Inter-facility Transfers** · Federal, state, county, and local regulations and guidance n Sv varies relative to inter-facility transfers LTCFs should limit transfers to Emergency Departments and hospitalizations to clinical conditions that require specialized testing and/or and acute or ICU level of care AMDA Clinical Practice Guidelines, the INTERACT program, and other similar tools should be used to help manage patients/residents in the facility whenever safe and feasible Patients/residents should have clearly documented advance directives if they are transferred to the extent that the patient/resident is capable of making their own decisions or there is a health care proxy available · Complete critical clinical information should accompany patients/residents using state mandated forms, the INTERACT transfer form, or a similar form that includes the information listed on the INTERACT transfer data list **FAU** MEDICINE

#### Key Points: Admissions to LTCFs from Hospitals

- Hospital patients should be discharged home whenever enough support is available to manage them safely outside of a post-acute facility
- Unless otherwise overridden by state, county or local regulations:
  - COVID-19 positive patients should no longer have symptoms and two negative tests 24 hours apart before being transferred from hospital to LTCF or meet CDC criteria
    - $\circ~$  No fever or respiratory symptoms for 72 hours and 7 or more days since onset of symptoms
  - Any patient being transferred from hospital to LTCF for any condition without a COVID-19 test result should be presumed to be infected, and isolated for at least 7 days
    - Based on risk of acquiring the virus in the hospital and nonspecificity of symptoms
- Hospitals should provide critical clinical information to post-acute settings using state information should accompany patients/residents using state mandated forms, the INTERACT transfer form, or a similar form that includes the information listed on the INTERACT transfer data list
  - This is especially important in settings where clinicians and other health professionals cannot access the hospital electronic medical record

## **FAU** MEDICINE



## COVID-19 in Long-Term Care Facilities: An Update

### Next Steps and Planning for the Future: Testing and Treatment

- As rapid testing and self-testing becomes more available, it will be easier to test all patients/residents and staff, and quarantine them as appropriate
- False negative tests do occur
- Testing serum will help identify who has been infected
  - o This will help with quarantine and staffing decisions
  - Convalescent serum/plasma may be a therapeutic option, however:
    - Not all people develop high antibody levels
    - · Duration of immunity is unknown may be a few months



#### Next Steps and Planning for the Future: Testing and Treatment

#### Currently there is no evidence-based drug treatment for COVID-19

- Hydroxychloroquine, with or without azithromycin may be helpful in treating the intense inflammatory response, but:
  - The data are basically anecdotal; results of controlled trials are pending but unlikely to include LTCF patients/residents
  - The drug has numerous potentially severe adverse effects, including sudden death in people with prolonged QT interval, and electrolyte and liver function abnormalities
  - Several potentially serious drug interactions
  - · If it is used:
  - · Consent should be documented
  - EKG performed before treatment
  - Guidance on dosing (intended for hospitals) was removed from the CDC website
- Other drugs, including antiviral agents and immune modulators are under investigation as is convalescent serum/plasma
- Vaccines are under development and should help prevent future waves of COVID disease

## **FAU** MEDICINE








# COVID-19 in Long-Term Care Facilities: An Update

#### Next Steps and Planning for the Future: Alternative Sites of Care

- Many areas are developing plans for alternative sites of care for patients who are suspected of or are recovering from COVID-19
  - Converting entire LTCFs
  - Using unoccupied wings of existing facilities
  - Critical access hospitals with swing beds
  - Temporary facilities
- Planning is complicated and requires cooperation between LTCFs, hospitals, county and state authorities
  - Regulatory, financial and liability issues need to be addressed
- Staffing and adequate PPE will be challenging







FLORIDA ATLANTIC UNIVERSITY		
Questions?	Comments?	Suggestions?
Joseph G. Ouslander, MD jousland@health.fau.edu		
FAU MEDICINE		